arXiv:math/0509689v2 [math.RA] 14 Nov 2005

Algebra / Algèbre

Graded algebras and multilinear forms

Michel Dubois-Violette [a]

[a] *Laboratoire de Physique Théorique, UMR 8627, Université Paris XI, Bâtiment 210, F-91 405 Orsay Cedex, France*

Abstract

We give a description of the connected graded algebras which are finitely generated and presented of global dimension 2 or 3 and which are Gorenstein. These algebras are constructed from multilinear forms. We generalize the construction by associating homogeneous algebras to multilinear forms. The homogeneous algebras which are Koszul of finite global dimension and which are Gorenstein are of this type.

Résumé

Algèbres graduées et formes multilinéaires. Nous donnons une description des algèbres graduées connexes de présentation finie et de dimension globale 2 ou 3 qui sont Gorenstein. Ces algèbres sont construites à partir de formes multilinéaires. Cette construction est généralisée en associant des algèbres homogènes aux formes multilinéaires. Sont de ce type les algèbres homogènes Koszul-Gorenstein de dimension globale finie.

Version française abrégée On s'intéresse aux algèbres graduées connexes de présentation finie sur un corps $\mathbb{K}$ algébriquement clos de caractéristique zéro. Une telle algèbre engendrée par un nombre fini d'éléments en degré 1 qui est de dimension globale $D = 2$ ou $D = 3$ et qui est Gorenstein est N-homogène et Koszul avec $N = 2$ pour $D = 2$ et $N \geq 2$ pour $D = 3$, [5]. Pour $D = 2$, on a le résultat suivant.*Soit B une forme bilinéaire non dégénérée sur $\mathbb{K}^q$ de composantes $B_{\mu\nu}$ dans la base canonique. L'algèbre quadratique $\mathcal{A}$ engendrée par les éléments x^λ ($\lambda \in \{1, \dots, q\}$) avec relation $B_{\mu\nu}x^\mu x^\nu = 0$ (on utilise partout la convention usuelle de sommation sur les indices répétés en haut et en bas) est Koszul de dimension globale 2 et Gorenstein. Inversement, toute algèbre quadratique engendrée par q éléments x^λ qui est Koszul de dimension globale 2 et Gorenstein est de ce type pour une forme bilinéaire non dégénérée B sur $\mathbb{K}^q$.* Nous aurons besoin des concepts suivants pour les formes multilinéaires. Soit V un espace vectoriel et n un entier avec $n \geq 1$; une forme $(n+1)$-linéaire W sur V sera dite *prérégulière* si elle vérifie les conditions (i) et (ii) suivantes :
(i) Si $X \in V$ est tel que $W(X, X_1, \dots, X_n) = 0$ pour tout $X_1, \dots, X_n \in V$ alors $X = 0$,

---

Email address: `Michel.Dubois-Violette@th.u-psud.fr` (Michel Dubois-Violette).

(ii) $\exists Q_W \in GL(V)$ tel que $W(X_0,\dots,X_{n-1},X_n)=W(Q_WX_n,X_0,\dots,X_{n-1})$ pour tout $X_0,\dots,X_n\in V$.
Soit N un entier avec $N\geq 2$; une forme $(N+1)$-linéaire sur V sera dite *3-régulière* si elle est prérégulière et vérifie la condition (iii) suivante :
(iii) Si $L_0,L_1\in \mathrm{End}(V)$ sont tels que $W(L_0X_0,X_1,X_2,\dots,X_N)=W(X_0,L_1X_1,X_2,\dots,X_N)$ pour tout $X_0,\dots,X_N\in V$ alors $L_0=L_1=\lambda\mathbb{1}$ avec $\lambda\in\mathbb{K}$.
Pour $D=3$, on a le résultat suivant. *Soit W une forme $(N+1)$-linéaire 3-régulière sur $\mathbb{K}^q$ de composantes $W_{\lambda_0\dots\lambda_N}$ dans la base canonique. L'algèbre N-homogène $\mathcal{A}$ engendrée par les éléments x^λ ($\lambda\in\{1,\dots,q\}$) avec relations $W_{\lambda\lambda_1\dots\lambda_N}x^{\lambda_1}\dots x^{\lambda_N}=0$ ($\lambda\in\{1,\dots,q\}$) est Koszul de dimension globale 3 et Gorenstein. Inversement, toute algèbre N-homogène engendrée par q éléments x^λ qui est Koszul de dimension globale 3 et Gorenstein est de ce type pour une forme $(N+1)$-linéaire 3-régulière W sur $\mathbb{K}^q$.* Cela suggère la construction générale suivante. Soient m et N deux entiers tels que $m\geq N\geq 2$ et soit $W\neq 0$ une forme m-linéaire sur $\mathbb{K}^q$, on définit l'algèbre N-homogène $\mathcal{A}(W,N)$ comme l'algèbre engendrée par q éléments x^λ avec relations $W_{\lambda_1\dots\lambda_{m-N}\mu_1\dots\mu_N}x^{\mu_1}\dots x^{\mu_N}=0$ ($\lambda,\lambda_i\in\{1,\dots,q\}$). Les résultats précédents admettent la généralisation partielle suivante. *Soit $\mathcal{A}$ une algèbre N-homogène qui est Koszul de dimension globale finie D et qui est Gorenstein ; on a $\mathcal{A}=\mathcal{A}(W,N)$ pour une forme m-linéaire prérégulière W sur $\mathbb{K}^q$, $q=dim(\mathcal{A}_1)$. Pour $N\geq 3$ on a $m=Np+1$ et $D=2p+1$ pour un entier $p\geq 1$ tandis que pour $N=2$ on a $m=D$.* La dernière partie de cet énoncé, qui implique que pour les dimensions globales paires les algèbres homogènes Koszul-Gorenstein sont quadratiques, est un cas particulier d'un résultat de [5] (proposition 5.3).

1. Introduction

One of our aims is to study the connected graded algebras which are finitely generated in degree 1 and finitely presented with relations of degrees ≥ 2 and which are of low global dimension D, $D=2$ and $D=3$. We further impose to these algebras to be Gorenstein. As pointed out in [5] (Proposition 5.2) such an algebra is N-homogeneous and Koszul with $N=2$ for $D=2$ and $N\geq 2$ for $D=3$. Our second more general objective is the study of the N-homogeneous algebras ($N\geq 2$) which are Koszul of arbitrary finite global dimension $D\geq 2$ and which are Gorenstein. For $D=2$, it is shown in Section 2 that these algebras are classified by the nondegenerate bilinear forms modulo the action of the linear group. For $D=3$, we show in Section 3 that these algebras are classified by nondegenerate $(N+1)$-linear forms satisfying a regularity condition called 3-regularity modulo the action of the linear group. In Section 4 we introduce and study homogeneous algebras associated with multilinear forms. It is pointed out that the Koszul homogeneous algebras of finite global dimension D which are Gorenstein belong to this class which generalizes the previous results for $D=2,3$.
Throughout the paper $\mathbb{K}$ denotes a field which is algebraically closed, of characteristic zero and all the algebras and vector spaces are over $\mathbb{K}$. In the following N, D and q denote integers greater than or equal to 2 and we use the Einstein summation convention of repeated up down indices in the formulas. For the notion of koszulity for homogeneous algebras introduced in [3] we refer to [3] and to [4]. For Koszul duality, Koszul N-complexes... for N-homogeneous algebras we refer to [4]; our notations are those of [4].

2. Global dimension $D=2$

As explained in the introduction, the connected graded algebras which are finitely generated in degree 1 and finitely presented of global dimension 2 and which are Gorenstein are the quadratic Koszul algebras of global dimension 2 which are Gorenstein. These algebras are characterized by the following theorem.

Theorem 2.1 *Let B be a nondegenerate bilinear form on $\mathbb{K}^q$ $(q\geq 2)$ with components $B_{\mu\nu}=B(e_\mu,e_\nu)$ in the canonical basis $(e_\lambda)_{\lambda\in\{1,\dots,q\}}$ of $\mathbb{K}^q$. Then the quadratic algebra $\mathcal{A}$ generated by the elements x^λ $(\lambda\in\{1,\dots,q\})$ with the relation $B_{\mu\nu}x^\mu x^\nu=0$ is Koszul of global dimension 2 and Gorenstein. Conversely, any quadratic algebra generated by q elements x^λ which is Koszul of global dimension 2 and Gorenstein is of the above kind for some non degenerate bilinear form B on $\mathbb{K}^q$. Two nondegenerate bilinear forms on $\mathbb{K}^q$ which are on the same $GL(q,\mathbb{K})$-orbit correspond to isomorphic algebras.*
Proof. Let $\mathcal{A}$ be the quadratic algebra generated by the x^λ with the relation $B_{\mu\nu}x^\mu x^\nu=0$. Then the dual quadratic algebra $\mathcal{A}^!$ is generated by elements $y_\lambda(\lambda\in\{1,\dots,q\})$ with relations $y_\mu y_\nu=(1/q)B_{\mu\nu}B^{\rho\tau}y_\tau y_\rho$ where $B^{\mu\nu}$ are the matrix elements of the inverse of the matrix $(B_{\mu\nu})$; i.e. $B_{\mu\lambda}B^{\lambda\nu}=\delta^\nu_\mu$. It follows that $\mathcal{A}^!_0=\mathbb{K}\mathbb{1}\simeq\mathbb{K}, \mathcal{A}^!_1=\oplus_\lambda\mathbb{K}y_\lambda\simeq\mathbb{K}^q, \mathcal{A}^!_2=\mathbb{K}B^{\beta\alpha}y_\alpha y_\beta\simeq\mathbb{K}$ and $\mathcal{A}^!_n=0$ for $n\geq 3$. The Koszul complex of $\mathcal{A}$ reads

$$0\rightarrow\mathcal{A}\xrightarrow{x^tB}\mathcal{A}^q\xrightarrow{x}\mathcal{A}\rightarrow 0 \tag{1}$$

where $\mathcal{A}^q=(\mathcal{A},\dots,\mathcal{A})$, x is right multiplication by the column (x^λ) while x^tB is right multiplication by the line $(x^\mu B_{\mu\nu})$. This complex is acyclic in degrees ≥ 1 so the algebra $\mathcal{A}$ is Koszul. The Gorenstein property follows by transposition and by using the invertibility of the matrix $(B_{\mu\nu})$.

Assume now that $\mathcal{A}$ is a quadratic algebra generated by elements x^λ $(\lambda\in\{1,\dots,q\})$ which is Koszul of global dimension 2 and Gorenstein. Then the Gorenstein property implies that the space of the quadratic relations of $\mathcal{A}$ is of dimension 1, i.e. that the relations of $\mathcal{A}$ read $B_{\mu\nu}x^\mu x^\nu=0$ for some non zero bilinear form B on $\mathbb{K}^q$. The Koszul complex reads again as (1) and the Gorenstein property implies that the matrix $(B_{\mu\nu})$ is invertible, i.e. that B is nondegenerate. The fact that $\mathcal{A}$ does only depend on the $GL(q,\mathbb{K})$-orbit of B is straightforward.□

For $q\geq 3$ the algebra has exponential growth while for $q=2$ it has polynomial growth so is regular in the sense of [1]. In the latter case it is easy to classify the $GL(2,\mathbb{K})$-orbits of nondegenerate bilinear forms on $\mathbb{K}^2$ according to the rank of their symmetric part [10] and one recovers the usual description of regular algebras of global dimension 2, [12], [1].

The algebra $\mathcal{A}$ of Theorem 2.1 corresponds to the natural quantum space for the action of the quantum group of the nondegenerate bilinear form defined in [10].

3. Global dimension $D=3$

In order to state the analog of Theorem 2.1 for homogeneous algebras which are Koszul of global dimension 3 and which are Gorenstein, we need the following concepts for multilinear forms.
Definition 3.1 *Let V be a vector space and n be an integer with $n\geq 1$. A $(n+1)$-linear form W on V will be said to be preregular iff it satisfies the following conditions (i), (ii).*
(i) If $X\in V$ is such that one has $W(X,X_1,\dots,X_n)=0$ for any $X_1,\dots,X_n\in V$, then $X=0$.
(ii) There is an invertible linear transformation $Q_W\in GL(V)$ such that one has

$$W(X_0,\dots,X_{n-1},X_n)=W(Q_WX_n,X_0,\dots,X_{n-1})$$

for any $X_0,\dots,X_n\in V$.

Let W be a preregular $(n+1)$-linear form on V. Then the $Q_W\in GL(V)$ such that (ii) is satisfied is unique and given any $p\in\{0,\dots,n\}$ the condition $W(X_1,\dots,X_p,X,X_{p+1},\dots,X_n)=0$ for any $X_1,\dots,X_n\in V$ implies $X=0$. Furthermore the space of these W is stable by the action $W\mapsto W^L$ of $GL(V)$ defined by $W^L(X_0,\dots,X_n)=W(L^{-1}X_0,\dots,L^{-1}X_n)$ and one has $Q_{W^L}=LQ_WL^{-1}$ for $L\in GL(V)$. Notice that a bilinear form is preregular iff it is nondegenerate.

Definition 3.2 *Let N be an integer with $N \geq 2$. A $(N+1)$-linear form W on V will be said to be 3-regular iff it is preregular and satisfies the following condition (iii).*
(iii) If $L_0, L_1 \in End(V)$ are such that one has $W(L_0X_0, X_1, X_2, \dots, X_N) = W(X_0, L_1X_1, X_2, \dots, X_N)$ for any $X_0, \dots, X_N \in V$, then $L_0 = L_1 = \lambda \mathbb{1}$ for some $\lambda \in \mathbb{K}$.
The subpace $\mathcal{R}^N_3(V)$ of all 3-regular $(N+1)$-linear forms on V is also stable by the action of $GL(V)$.
Theorem 3.3 *Let W be a 3-regular $(N+1)$-linear form on $\mathbb{K}^q$ with components $W_{\lambda_0\dots\lambda_N} = = W(e_{\lambda_0}, \dots, e_{\lambda_N})$ in the canonical basis $(e_\lambda)_{\lambda\in\{1,\dots,q\}}$ of $\mathbb{K}^q$. Then the N-homogeneous algebra $\mathcal{A}$ generated by elements $x^\lambda (\lambda \in \{1, \dots, q\})$ with relations $W_{\lambda\lambda_1\dots\lambda_N} x^{\lambda_1} \dots x^{\lambda_N} = 0$ ($\lambda \in \{1, \dots, q\}$) is Koszul of global dimension 3 and Gorenstein. Conversely, any N-homogeneous algebra generated by q elements x^λ which is Koszul of global dimension 3 and Gorenstein is of the above kind for some 3-regular $(N+1)$-linear form W on $\mathbb{K}^q$. Two 3-regular $(N+1)$-linear forms on $\mathbb{K}^q$ which are on the same $GL(q,\mathbb{K})$-orbit correspond to isomorphic algebras.*
Proof. Let $\mathcal{A}$ be the N-homogeneous algebra generated by the x^λ with relation $W_{\lambda\lambda_1\dots\lambda_N} x^{\lambda_1} \dots x^{\lambda_N} = 0$. Then one has the complex

$$0 \to \mathcal{A}\otimes \underline{W} \xrightarrow{d} \mathcal{A}\otimes R \xrightarrow{d^{N-1}} \mathcal{A}\otimes E \xrightarrow{d} \mathcal{A} \to \mathbb{K} \to 0 \tag{2}$$

where $E = \oplus_\lambda \mathbb{K}x^\lambda$, $R = \oplus_\lambda \mathbb{K} W_{\lambda\lambda_1\dots\lambda_N} x^{\lambda_1} \otimes \dots \otimes x^{\lambda_N} \subset E^{\otimes^N}$, $\underline{W}$ is the one-dimensional subspace $\mathbb{K}W_{\lambda_0\dots\lambda_N} x^{\lambda_0} \otimes \dots x^{\lambda_N}$ of $E^{\otimes^{N+1}}$ and where d is induced by $a\otimes(x^0\otimes x^1\otimes\dots\otimes x^n) \mapsto ax^0\otimes(x^1\otimes\dots\otimes x^n)$ of $\mathcal{A}\otimes E^{\otimes^{n+1}}$ into $\mathcal{A}\otimes E^{\otimes^n}$ (the arrow $\mathcal{A} \to \mathbb{K}$ being induced by the projection onto the degree 0). By the very definition of $\mathcal{A}$ by generators and relations, the sequence (2) is exact at the first 3 terms from the right, exactness at $\mathcal{A}\otimes\underline{W}$ follows from preregularity while exactness at $\mathcal{A}\otimes R$ is equivalent to (iii) once preregularity is assumed. Thus Sequence (2) is exact, it is a free resolution of the trivial left $\mathcal{A}$-module $\mathbb{K}$. It is easy to see then that $\underline{W} = E\otimes R \cap R \otimes E$ so $\mathcal{A}$ is Koszul of global dimension 3 and (2) is the corresponding Koszul resolution. The Gorenstein property follows from (ii) with invertibility of Q_W.
Conversely given a N-homogeneous algebra $\mathcal{A}$ which is Koszul of global dimension 3 and Gorenstein, the Koszul resolution of the trivial left $\mathcal{A}$-module $\mathbb{K}$ reads

$$0 \to \mathcal{A}\otimes(E\otimes R\cap R\otimes E) \xrightarrow{d} \mathcal{A}\otimes R \xrightarrow{d^{N-1}} \mathcal{A}\otimes E \xrightarrow{d} \mathcal{A} \to \mathbb{K} \to 0$$

and Gorenstein property implies $\dim(E\otimes R\cap R\otimes E) = 1$ so $E\otimes R\cap R\otimes E = \mathbb{K}W$ for some $W \in E^{\otimes^{N+1}}$ which (again via Gorenstein property) must satisfy conditions (i) and (ii) and therefore also (iii).

The fact that $\mathcal{A}$ does only depend on the $GL(q,\mathbb{K})$-orbit of W is straightforward. $\square$

The Poincaré series of such a N-homogeneous algebra $\mathcal{A}$ which is Koszul of global dimension 3 and which is Gorenstein is given by [2], [11] $P_\mathcal{A}(t) = (1 - qt + qt^N - t^{N+1})^{-1}$ where $q = \dim(\mathcal{A}_1)$ as before. It follows that $\mathcal{A}$ has exponential growth if $q + N > 5$. The case $q = 2$ and $N = 2$ is impossible so it remains the cases $q = 3, N = 2$ and $q = 2, N = 3$ for which one has polynomial growth [1]. These latter cases are the object of [1] and one encounters there various values of Q_W in $GL(3,\mathbb{K})$ and in $GL(2,\mathbb{K})$. Examples with $N = 3$ and arbitrary values of q are the Yang-Mills algebra [7] and its deformations [9] for which $Q_W = \mathbb{1}$ and the super Yang-Mills algebra and its deformations [9] for which $Q_W = -\mathbb{1}$.

4. Homogeneous algebras and multilinear forms

Let m and N be two integers such that $m \geq N \geq 2$ and let W be a m-linear form on $\mathbb{K}^q$ with $W \neq 0$. We denote by $\underline{W}$ the 1-dimensional space of multilinear forms spanned by W and let $W_{\lambda_1\dots\lambda_m} = W(e_{\lambda_1}, \dots, e_{\lambda_m})$ be the components of W in the canonical basis $(e_\lambda)_{\lambda\in\{1,\dots,q\}}$ of $\mathbb{K}^q$. Let $\mathcal{A} = \mathcal{A}(W,N)$ be the N-homogeneous algebra generated by elements x^λ with relations $W_{\lambda_1\dots\lambda_{m-N}\mu_1\dots\mu_N} x^{\mu_1} \dots x^{\mu_N} = 0$

$(\lambda, \lambda_i \in \{1, \dots, q\})$, in other words $\mathcal{A}(W,N) = A(E,R)$ with $E = \oplus_\lambda \mathbb{K}x^\lambda$ and with space of relations $R = \sum_{\lambda_i} \mathbb{K}W_{\lambda_1\dots\lambda_{m-N}\mu_1\dots\mu_N} x^{\mu_1} \otimes \dots \otimes x^{\mu_N} \subset E^{\otimes^N}$. The vector space E can be interpreted as the dual of $\mathbb{K}^q$ and its basis (x^λ) as the dual basis of (e_λ) while $\underline{W}$ is the 1-dimensional subspace of $E^{\otimes^m}$ spanned by $W \in E^{\otimes^m}$. To $\underline{W} \subset E^{\otimes^m}$ we associate the family of subspaces $\underline{W}^{(n)} \subset E^{\otimes^{m-n}}$, $m \geq n \geq 0$, defined by $\underline{W}^{(0)} = \underline{W}$ and $\underline{W}^{(n)} = \sum_{\lambda_i} \mathbb{K}W_{\lambda_1\dots\lambda_n\mu_1\dots\mu_{m-n}} x^{\mu_1} \otimes \dots \otimes x^{\mu_{m-n}}$ for $m \geq n \geq 1$. Consider the sequence

$$0 \to \mathcal{W}_m \xrightarrow{d} \mathcal{W}_{m-1} \xrightarrow{d} \dots \xrightarrow{d} \mathcal{W}_2 \xrightarrow{d} \mathcal{W}_1 \xrightarrow{d} \mathcal{W}_0 \to 0 \tag{3}$$

of free left $\mathcal{A}$-modules and $\mathcal{A}$-module-homomorphisms with $\mathcal{W}_n \subset \mathcal{A} \otimes E^{\otimes^n}$ defined by

$$\mathcal{W}_n = \begin{cases} \mathcal{A} \otimes \underline{W}^{(m-n)} & \text{if } m \geq n \geq N \\ \mathcal{A} \otimes E^{\otimes^n} & \text{if } N-1 \geq n \geq 0 \end{cases}$$

and where the homomorphisms d are induced by the homomorphisms of $\mathcal{A} \otimes E^{\otimes^{n+1}}$ into $\mathcal{A} \otimes E^{\otimes^n}$ defined by $a \otimes (v_0 \otimes v_1 \otimes \dots v_n) \mapsto av_0 \otimes (v_1 \otimes \dots \otimes v_n)$ for $m > n \geq 0$, $a \in \mathcal{A}$ and $v_i \in E$ $(= \mathcal{A}_1)$.

Proposition 4.1 *Assume that the m-linear form W is preregular. Then Sequence (3) is a sub-N-complex of the Koszul N-complex $K(\mathcal{A})$.*

Proof. One has $W^{(m-n)} \subset E^{\otimes^{n-N}} \otimes R$ for $m \geq n \geq N$. By Property (ii) of Definition 3.1, relations $W_{\lambda_a\dots\lambda_{m-N}\mu_1\dots\mu_N} x^{\mu_1} \dots x^{\mu_N} = 0$ are equivalent to relations $W_{\lambda_{r+1}\dots\lambda_{m-N}\mu_1\dots\mu_N\nu_1\dots\nu_r} x^{\mu_1} \dots x^{\mu_N} = 0$ for $m-N \geq r \geq 0$. It follows that $W^{(m-n)} \subset E^{\otimes^{n-N-r}} \otimes R \otimes E^{\otimes^r}$ so $\mathcal{W}_n \subset \mathcal{A} \otimes \cap_r E^{\otimes^{n-N-r}} \otimes R \otimes E^{\otimes^r} = K_n(\mathcal{A})$ for $m \geq n \geq N$ and one has the result. □

In the remaining part of this section W is assumed to be preregular, Q_W denotes the corresponding element of $GL(q,\mathbb{K}) = GL(E^*)$ and $\mathcal{A} = \mathcal{A}(W,N) = A(E,R)$ is the N-homogeneous algebra defined above. One has $W \in \cap_r E^{m-N-r} \otimes R \otimes E^r = \mathcal{A}^{!*}_m$ so W composed with the canonical projection $\mathcal{A}^! \to \mathcal{A}^!_m$ on degree m defines a linear form ω_W on $\mathcal{A}^!$.

Proposition 4.2 *Let $W, \mathcal{A}, \mathcal{A}^!, Q_W, \omega_W$ be as above.*
(i) $Q_W \in GL(E^*)$ induces an automorphism σ_W of the N-homogeneous algebra $\mathcal{A}^!$.
(ii) One has $\omega_W(xy) = \omega_W(\sigma_W(y)x)$ for $x, y \in \mathcal{A}^!$.

Proof. Q_W induces an automorphism $\tilde{\sigma}_W$ of degree zero of the tensor algebra $T(E^*)$. Let $\tilde{x}$ be in $R^\perp$ i.e. $\tilde{x} = \rho^{\mu_1\dots\mu_N} e_{\mu_1} \otimes \dots \otimes e_{\mu_N}$ such that $W_{\lambda_1\dots\lambda_{m-N}\mu_1\dots\mu_N} \rho^{\mu_1\dots\mu_N} = 0$, $\forall \lambda_i$. Then since $W \circ Q_W^{\otimes^m} = W$ one has $Q^{\rho_1}_{\lambda_1} \dots Q^{\rho_{m-N}}_{\lambda_{m-N}} W_{\rho_1\dots\rho_{m-N}\nu_1\dots\nu_N} Q^{\nu_1}_{\mu_1} \dots Q^{\nu_N}_{\mu_N} \rho^{\mu_1\dots\mu_N} = 0$ i.e. $W_{\lambda_1\dots\lambda_{m-N}\nu_1\dots\nu_N} Q^{\nu_1}_{\mu_1} \dots Q^{\nu_N}_{\mu_N} \rho^{\mu_1\dots\mu_N} = 0$ $(\forall \lambda_i)$ which means that $\tilde{\sigma}_W(\tilde{x})$ is in $R^\perp$. Thus $\tilde{\sigma}_W$ passes to the quotient and defines an automorphism σ_W of $\mathcal{A}^!$ which proves (i). (ii) is then obvious in view of (ii) in Definition 3.1. □

Theorem 4.3 *Let $\mathcal{A}$ be a N-homogeneous Koszul algebra of finite global dimension D which is Gorenstein. Then $\mathcal{A} = \mathcal{A}(W,N)$ for some preregular m-linear form on $\mathbb{K}^q$, $q = dim(\mathcal{A}_1)$. Furthermore if $N \geq 3$ then $m = Np+1$ and $D = 2p+1$ for some integer $p \geq 1$, while for $N = 2$ one has $m = D$.*

Proof. The Koszul resolution starts as $\dots \xrightarrow{d} \mathcal{A} \otimes \mathcal{A}^{!*}_N \xrightarrow{d^{N-1}} \mathcal{A} \otimes \mathcal{A}_1 \xrightarrow{d} \mathcal{A} \to \mathbb{K} \to 0$ and must end as $0 \to \mathcal{A} \otimes \mathcal{A}^{!*}_m \xrightarrow{d} \mathcal{A} \otimes \mathcal{A}^{!*}_{m-1} \xrightarrow{d^{N-1}} \dots$ in view of the Gorenstein property. This implies either $N = 2$ and $m = D$ or if $N > 2$, $m = Np+1$ and $D = 2p+1$ for some integer $p \geq 1$. The Gorenstein property implies also that $\dim(\mathcal{A}^{!*}_m) = 1$ and $\dim(\mathcal{A}^{!*}_{m-1}) = q$. Let W be a generator of $\mathcal{A}^{!*}_m \subset \mathcal{A}_1^{\otimes^m}$. Identifying $\mathcal{A}^*_1$ with $\mathbb{K}^q$, W is a m-linear form on $\mathbb{K}^q$, $\mathcal{A}^{!*}_{m-1} = \underline{W}^{(1)}$ and W satisfies Condition (i) of Definition 3.1. Theorem 5.4 of [5] implies then that $\mathcal{A}^{!*}_n = \underline{W}^{(m-n)}$ for $n = Nk$ and $n = Nk+1$, $k \in \mathbb{N}$. So, in particular the space of relations $R = \mathcal{A}^{!*}_N$ coincides with $\underline{W}^{(m-N)}$ which means that $\mathcal{A} = \mathcal{A}(W,N)$. Condition (ii) of Definition 3.1 follows also (see e.g. Corollary 5.12 of [5]), so W is preregular.□

As example with $N = 2$ and $m = D = 4$, consider the algebra $\mathcal{A}_{\mathbf{u}}$ of [6] and [8]. This is the quadratic algebra generated by 4 elements x^λ, $\lambda \in \{0,1,2,3\}$, with relations

$$\cos(\varphi_0-\varphi_k)[x^0,x^k]=i\sin(\varphi_\ell-\varphi_m)\{x^\ell,x^m\}\ \text{and}\ \cos(\varphi_\ell-\varphi_m)[x^\ell,x^m]=i\sin(\varphi_0-\varphi_k)\{x^0,x^k\}$$

for any cyclic permutation (k,ℓ,m) of (1,2,3) and where $\{A,B\}=AB+BA$. This algebra is Koszul of global dimension 4 and Gorenstein whenever none of these six relations becomes trivial and then one has as explained in [6], the nontrivial Hochschild cycle

$$\begin{aligned}W=\tilde{\mathrm{ch}}_{\frac{3}{2}}(U_{\mathbf{u}})=&-\sum_{\alpha,\beta,\gamma,\delta}\epsilon_{\alpha\beta\gamma\delta}\cos(\varphi_\alpha-\varphi_\beta+\varphi_\gamma-\varphi_\delta)x^\alpha\otimes x^\beta\otimes x^\gamma\otimes x^\delta\\&+i\sum_{\mu,\nu}\sin(2(\varphi_\mu-\varphi_\nu))x^\mu\otimes x^\nu\otimes x^\mu\otimes x^\nu\end{aligned}$$

which may be considered as a 4-linear form on $\mathbb{K}^4$. One checks that one has $\mathcal{A}_{\mathbf{u}}=\mathcal{A}(W,2)$ and that $Q_W=-\mathbb{1}$ here.

References

[1] M. Artin, W.F. Schelter. Graded algebras of global dimension 3. *Adv. Math.* 66:171–216, 1987.

[2] M. Artin, J. Tate, M. Van den Bergh. Modules over regular algebras of dimension 3. *Invent. Math.*, 106:335–388, 1991.

[3] R. Berger. Koszulity for nonquadratic algebras. *J. Algebra* 239:705–734, 2001.

[4] R. Berger, M. Dubois-Violette, M. Wambst. Homogeneous algebras. *J. Algebra* 261:172–185, 2003.

[5] R. Berger, N. Marconnet. Koszul and Gorenstein properties for homogeneous algebras. math.QA/0310070.

[6] A. Connes, M. Dubois-Violette. Noncommutative finite-dimensional manifolds. I.Spherical manifolds and related examples. *Commun. Math. Phys.* 230:539–579, 2002.

[7] A. Connes, M. Dubois-Violette. Yang-Mills algebra. *Lett. Math. Phys.* 61:149–158, 2002.

[8] A. Connes, M. Dubois-Violette. Moduli space and structure of noncommutative 3-spheres. *Lett. Math. Phys.* 66:91-121, 2003.

[9] A. Connes, M. Dubois-Violette. Yang-Mills and some related algebras. math-ph/0411062.

[10] M. Dubois-Violette, G. Launer. The quantum group of a non-degenerated bilinear form. *Phys. Lett.*245B:175-177, 1990.

[11] M. Dubois-Violette, T. Popov. Homogeneous algebras, statistics and combinatorics. *Lett. Math. Phys.* 61:159-170, 2002.

[12] R.S. Irving. Prime ideals of Ore extensions. *J. Algebra* 58:399–423, 1979.